

**Supplementary Table S7.** Latent effects of early-life exposure to DCA on the incidence of preneoplastic hepatocellular foci.

Duration of DCA exposure   control dH <sub>2</sub> O treatment (weeks)	0   93	4   89	10   83	26   67	52   41	93   0
<b>No. Evaluated (&lt;31 weeks)</b>	<b>30</b>	<b>0</b>	<b>30</b>	<b>20</b>	<b>0</b>	<b>0</b>
None observed						
<b>No. Evaluated (≥31 weeks)</b>	<b>60</b>	<b>23</b>	<b>61</b>	<b>59</b>	<b>52</b>	<b>33</b>
No. indiv (%)						
Focus, Any	3 (5)	5 (22)	8 (13)	11 (19)	14 (27)*	3 (9)
Basophilic Focus	1 (2)	3 (13)	8 (13)	11 (19)*	10 (19)*	0 (0)
Eosinophilic Focus	1 (2)	1 (4)	1 (2)	0 (0)	3 (6)	2 (6)
Clear/Vacuolated Cell Focus	1 (2)	2 (9)	1 (2)	1 (2)	1 (2)	1 (3)

Asterisk (\*) indicates significant difference (adjusted  $P < 0.05$ ) compared to control group by one-sided Fisher's Exact Test. Counts include early death cases evaluated by histopathology. First hepatocellular focus was observed at 31 weeks.